



Cite this article: Ottoni C, Rasteiro R, Willet R, Claeys J, Talloen P, Van de Vijver K, Chikhi L, Poblome J, Decorte R. 2016 Comparing maternal genetic variation across two millennia reveals the demographic history of an ancient human population in southwest Turkey. *R. Soc. open sci.* **3**: 150250. <http://dx.doi.org/10.1098/rsos.150250>

Received: 8 June 2015

Accepted: 20 January 2016

Subject Category:

Genetics

Subject Areas:

genetics

Keywords:

ancient DNA, approximate Bayesian computation, Turkey, Roman, Byzantine

Author for correspondence:

Claudio Ottoni

e-mail: claudio.ottoni@med.kuleuven.be

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsos.150250> or via <http://rsos.royalsocietypublishing.org>.

Comparing maternal genetic variation across two millennia reveals the demographic history of an ancient human population in southwest Turkey

Claudio Ottoni^{1,2,4}, Rita Rasteiro⁵, Rinse Willet⁶, Johan Claeys³, Peter Talloen³, Katrien Van de Vijver¹, Lounès Chikhi^{7,8,9}, Jeroen Poblome³ and Ronny Decorte^{2,4}

¹Center for Archaeological Sciences, ²Department of Imaging & Pathology, and

³Sagalassos Archaeological Research Project, University of Leuven, Leuven, Belgium

⁴Laboratory of Forensic Genetics and Molecular Archaeology, UZ Leuven, Belgium


⁵Department of Genetics, School of History, University of Leicester, Leicester, UK

⁶Department of Humanities, Institute of History, Leiden University, Leiden, The Netherlands

⁷CNRS, Université Paul Sabatier, ENFA; UMR5174 EDB (Laboratoire Évolution & Diversité Biologique), Toulouse, France

⁸Université Toulouse 3 Paul Sabatier, CNRS; UMR5174 EDB, Toulouse, France

⁹Instituto Gulbenkian de Ciência, Oeiras, Portugal

 RR, 0000-0002-4217-3060

More than two decades of archaeological research at the site of Sagalassos, in southwest Turkey, resulted in the study of the former urban settlement in all its features. Originally settled in late Classical/early Hellenistic times, possibly from the later fifth century BCE onwards, the city of Sagalassos and its surrounding territory saw empires come and go. The Plague of Justinian in the sixth century CE, which is considered to have caused the death of up to a third of the population in Anatolia, and an earthquake in the seventh century CE, which is attested to have devastated many monuments in the city, may have severely affected the contemporary Sagalassos community. Human occupation continued, however, and Byzantine Sagalassos was eventually abandoned around 1200 CE. In order to investigate whether these historical events

resulted in demographic changes across time, we compared the mitochondrial DNA variation of two population samples from Sagalassos (Roman and Middle Byzantine) and a modern sample from the nearby town of Ağlasun. Our analyses revealed no genetic discontinuity across two millennia in the region and Bayesian coalescence-based simulations indicated that a major population decline in the area coincided with the final abandonment of Sagalassos, rather than with the Plague of Justinian or the mentioned earthquake.

1. Background

In the last three decades, palaeogenetics, the study of ancient DNA (aDNA), has proven invaluable to disentangle the biological and cultural processes in human (pre)history. The time-depth provided by the analysis of genetic variation in past populations is key to circumventing some of the limitations of interpreting present-day genetic patterns alone. In particular, the study of the maternally inherited mitochondrial DNA (mtDNA) has contributed to reveal past demographic scenarios previously undetected in ancient human communities.

Within the Eurasian continent, most aDNA studies on human populations have addressed various levels of temporal and geographical resolution, from wide-scale prehistoric population dynamics [1,2] to smaller-scale demographic events in more recent historical times [3,4].

Across multiple temporal layers of the history of our species, the importance of the Anatolian peninsula as a bridge between continents and cultures is unquestionable. The remains of the ancient city of sagalassos are situated in southwest Turkey, at an altitude of 1450–1600 m above sea level in the Taurus mountain range. Located 7 km north of the town of Ağlasun, in the Turkish province of Burdur (electronic supplementary material, figure S1), Sagalassos played a key role in the past of the wider region then known as Pisidia, as almost 30 years of archaeological research have revealed [5].

The first indications of human presence in the Ağlasun Valley can be placed at approximately 10 000 BCE in Epi-Palaeolithic times. The study region, which forms part of the Turkish Lake District, contains a range of well-studied Late Neolithic and Bronze Age sites [6]. The so-called Beysehir occupation phase (BO-phase) introduced a period of increased anthropogenic activity from around 1000 BCE, although the start date can vary quite significantly between specific areas. A hierarchical settlement landscape emerged during the Iron Age in the wider region. Finally, in late Classical/early Hellenistic times (possibly from the later fifth century BCE onwards), Sagalassos and Düzen Tepe—an adjacent settlement that was inhabited into the second century BCE—originated together with a group of small farms in the surrounding valley. From its origins up to its abandonment in the early thirteenth century CE, Sagalassos formed part of different empires administering Pisidia, namely that of the Achaemenids, the Hellenistic Successor Kings, the Roman and Byzantine Empires, and possibly their regional Seljuk sequel [7].

During the *Pax Romana*, a period of peace and reduced military engagement spanning more than two centuries from 25 BCE up to 180 CE, Sagalassos experienced one of its most flourishing periods. This is testified by an intense pottery production, large-scale exchange [8] and expansive urbanization into the early third century (electronic supplementary material, text). Sagalassos remained a densely occupied urban centre at least until the mid-sixth century. Afterwards, the urban framework changed, including the abandonment of some quarters in the town.

Two main factors are considered to have contributed to the traditional scenario of late antique (demographic) decline and the overall restructuring of Sagalassos' community. One factor is the first recorded outbreak of plague—the Plague of Justinian—which between 541 and 543 CE spread into Europe through the Mediterranean Basin [9], causing the death of up to a third of the population in Anatolia and the Eastern Mediterranean [10]. Pisidia may have suffered the same fate, even though no clear evidence of the spread of the plague up to Sagalassos and its wider region is so far present.

The second factor is seismic activity in the area. A severe earthquake destroyed several monuments in the urban centre in the course of the first half of the seventh century CE [11]. The community of Sagalassos lived on, albeit in different conditions, possibly resembling the larger villages that the Roman emperor Justinian associated with the region of Pisidia [12].

Whereas the demographic trajectories of the following centuries remain quite elusive, at least based on the current archaeological and historical evidence, the final abandonment of Sagalassos can be dated to the early thirteenth century CE, when a fortified settlement or *kastron* was destroyed, and the local population is considered to have shifted towards Ağlasun. By that time, Ağlasun was occupied by the Seljuk Turks [13] and is assumed to have been continuously inhabited up to the present time.

To better elucidate the demographic changes across time that affected the people living in Sagalassos and its territory, we have analysed the mtDNA variation in 24 individuals dated to Roman and Early Byzantine times, ranging from the early second up to the sixth century CE. The mtDNA sequences obtained were compared with a Middle Byzantine sample from Sagalassos previously studied [14] and with modern mtDNA sequences from the town of Ağlasun [15] (electronic supplementary material, table S2), which nowadays accounts for about 4100 inhabitants (<http://www.tursaga.com/>). Finally, we applied Bayesian coalescence-based approaches to identify the most plausible demographic scenarios supported by available DNA, historical and archaeological evidence.

2. Methods

2.1. Genetic analyses

Bone and teeth samples of 44 individuals dated to late Hellenistic (late first century BCE), Roman Imperial (first–third centuries CE), Late Roman (fourth–fifth centuries CE) and Early Byzantine times (sixth–seventh centuries CE), as determined by direct accelerator mass spectrometry radiocarbon dating, stratigraphic associations and contextual archaeological evidence, were available for aDNA analyses (electronic supplementary material, table S1).

Analyses were carried out in dedicated aDNA facilities of the Forensic Genetics and Molecular Archaeology Department of the University of Leuven (Belgium). The two hypervariable segments (HVS-I and HVS-II) of the mtDNA and eight diagnostic SNPs of the mtDNA coding region were amplified and sequenced in multiple independent experiments as described elsewhere [14]. Further details regarding methods, precautions taken to minimize contamination, and the criteria adopted to authenticate the aDNA sequences are described in the electronic supplementary material, text.

2.2. Data analyses

To unravel the genetic affinities of two ancient population samples from Sagalassos (Roman and Middle Byzantine) and the modern sample from Ağlasun to present-day populations, we generated an in-house database with more than 11 000 concatenated HVS-1 and HVS-2 mtDNA sequences of contemporary populations from the literature (electronic supplementary material, table S7). We used ARLEQUIN v. 3.5 [16] to compute pairwise F_{ST} genetic distances. The Slatkin's linearized F_{ST} values were represented in a two-dimensional multidimensional scaling plot with STATISTICA v. 8 (Statsoft) and for Sagalassos and Ağlasun they were plotted on a geographical map with SURFER v. 6 (Golden Software) using the Kriging method. The relative frequencies of haplotypes from Sagalassos and Ağlasun (excluded CRS haplotypes) shared with the modern populations were estimated with ARLEQUIN and displayed on a geographical map with SURFER. A median-joining network [17] of concatenated HVS1–HVS2 haplotypes from Sagalassos and Ağlasun was created with NETWORK v. 4.6 (www.fluxus-engineering.com).

To include a larger number of populations and ancient samples from the literature, a second database of more than 17 000 HVS-1 sequences was used for F_{ST} genetic distances analyses (electronic supplementary material, table S8).

Haplogroup frequencies from more than 180 ancient and modern populations were used for a principal component analysis (PCA) and represented in a two-dimensional plot with STATISTICA (electronic supplementary material, table S9). Detailed information about the statistical methods and the databases is given in the electronic supplementary material.

2.3. Simulation analyses

In order to identify the major events that took place during the history of Sagalassos, we first built demographic models (figure 1) involving either no change in population size or a population size change during one of the known catastrophic events that took place (plague, earthquake, abandonment). To compare the different models and identify those that best explain the data, we used an approximate Bayesian computation (ABC) framework [18,19] and implemented it together with the Bayesian serial SIMCOAL software [20,21], a program that simulates ancient and modern mtDNA data with sample sizes and dates identical to the real data. We also applied two different methods to test for continuity versus discontinuity between the two ancient samples from Sagalassos (Roman and Middle Byzantine) and Ağlasun (electronic supplementary material, text and figure S8).




Figure 1. Demographic scenarios used in the ABC analysis and their posterior probabilities. Two sets of different demographic scenarios were tested using ancient and modern mtDNA. In panel (a), three simple scenarios of population size change were compared to study the impact of a single past population contraction following the Plague of Justinian (PM), the seventh century earthquake (QM) and the abandonment (AM) of the city of Sagalassos. These scenarios assume a single constant female effective population size N_s , prior to the contraction, sampled from an ancestral female population of constant size N_{anc} , corresponding to the initial settlers of the region. A scenario of no size change (CM) since the BO-phase in the region until the present in Ağlasun was also tested as a null hypothesis. The posterior probabilities identified the AM scenario as the best model. In panel (b), scenarios with a second contraction added to the AM scenario were studied. This population size contraction corresponds to a size reduction during the plague and/or earthquake at time T_{pq} and was followed by either a period of recovery (PQR + AM) or constant population size (PQ + AM) until the abandonment of the city of Sagalassos. The parameter p_i (p_p , p_q and p_a for plague, quake and abandonment of the city, respectively) represents the proportion used to calculate the effective population size after the contraction (N_c) at time i , such that $N_c = p_i \times N_s$ for scenarios AM, PM and QM. In the more complex scenarios, N_c after the abandonment of the city is $p_{pq} \times p_a \times N_s$ (p_{pq} being the contraction proportion at time T_{pq}) and $p_a \times N_s$ for PQ + AM and PQR + AM, respectively. Both in (a) and (b), the posterior probabilities under each model are represented, calculated using the multinomial logistic regression (MLR) method of Beaumont [18], under an ABC framework.

Further details regarding the ABC validation, continuity tests and additional simulations accounting for potential maternal relationships in the Roman sample are fully described in the electronic supplementary material.

3. Results

3.1. Mitochondrial DNA genetic pool in Sagalassos across time

Authentic aDNA sequences were obtained for 22 Roman and two Early Byzantine individuals (electronic supplementary material, table S2), which were pooled in one population sample addressed here as Roman. Haplogroup frequencies in Roman and Middle Byzantine Sagalassos and Ağlasun (table 1) show overlapping 95% Bayesian credible intervals except for macro-haplogroup M in Middle Byzantine Sagalassos and Ağlasun.

The high diversity of mtDNA lineages observed in Sagalassos and Ağlasun is exhibited by the network (electronic supplementary material, figure S2). In several instances, haplotypes of the three samples (haplogroups W, K1, T2, U5a, H) differ by one or two mutational steps.

When compared with haplotypes of the database, Sagalassos shows a lower fraction of unique haplotypes (43% and 41%, respectively, in the Roman and Middle Byzantine samples) than Ağlasun (57%) and overall the haplotypes shared with present-day populations have a West Eurasian distribution (electronic supplementary material, figure S3 and tables S5–S6).

Figure 2 represents the pairwise genetic distances calculated from the database of HVS-1 and HVS-2 sequences on a geographical map. A typical West Eurasian variation in Sagalassos during Roman times is evident (figure 2a) and the F_{ST} values indicate in particular higher affinity with populations from the Caucasus, Southwest Asia, the Eastern Mediterranean, Southern Italy and some countries of continental Europe (Germany, Northern France; electronic supplementary material, table S3 and figure S4a). Northern Asian populations exhibited the largest values. A similar pattern of genetic affinities is observed in Middle Byzantine Sagalassos and in Ağlasun (figure 2b,c).

A Southwest Asian affinity of Sagalassos and Ağlasun is also detected when contrasting genetic distances with a larger number of populations at the level of HVS-1 variation (electronic supplementary material, table S4 and figures S4b and S6). Ancient Mesolithic, Neolithic and Bronze Age samples from Europe and Asia were included in the analysis and some from Early–Middle Neolithic central Europe [1] showed proximity with Sagalassos.

Table 1. Absolute and relative frequencies of mtDNA haplogroups. This table presents the estimated frequencies together with the 95% Bayesian credible intervals (CI), in Roman and Middle Byzantine Sagalassos and in the modern town of Ağlasun.

	Roman Sagalassos		Middle Byzantine Sagalassos ^a		Ağlasun ^b	
	% (N)	CI	% (N)	CI	% (N)	CI
R0/HV/H	17 (4)	7–36	29 (15)	19–43	25 (13)	15–38
I	0	0–14	0	0–7	2 (1)	0–10
N1	0	0–14	12 (6)	6–23	0	0–7
J	0	0–14	12 (6)	6–23	8 (4)	3–18
T	17 (4)	7–36	10 (5)	4–21	17 (9)	9–29
U1	4 (1)	1–20	4 (2)	1–13	0	0–7
U2	0	0–14	0	0–7	6 (3)	2–15
U3	0	0–14	6 (3)	2–16	0	0–7
U4	4 (1)	1–20	0	0–7	0	0–7
U5	4 (1)	1–20	2 (1)	0–10	4 (2)	1–13
U6	0	0–14	4 (2)	1–13	0	0–7
U7	4 (1)	1–20	0	0–7	0	0–7
U8 (incl. K)	29 (7)	15–49	12 (6)	6–23	13 (7)	7–25
W	4 (1)	1–20	8 (4)	3–19	2 (1)	0–10
X	17 (4)	7–36	2 (1)	0–10	0	0–7
M (incl. C)	0	0–14	0	0–7	15 (8)	8–27
?	0	0–14	0	0–7	9 (5)	4–20
Total (N)	24		51		53	

^aOttoni *et al.* [14].^bJehaes [15].

Concordant with the F_{ST} analysis, the haplogroup frequencies PCA (electronic supplementary material, figure S7) reveals affinity between ancient Sagalassos and Southwest Asian populations, and an even closer relationship with samples from Caucasus, Southern Italy (Sicily, Apulia, Campania) and the ancient samples of Early and Middle Neolithic central European farmers and Bronze Age Crete.

3.2. Identifying, quantifying and dating population size changes

Figure 1 represents the two sets of demographic scenarios tested together with their posterior probabilities [22]. These scenarios assume continuity between the Sagalassos and Ağlasun samples, and this continuity was tested and not rejected (electronic supplementary material, text and figure S8). In figure 1*a*, the model with the highest support is the AM (91%), where the population contraction corresponding to the abandonment of the city of Sagalassos is very strong (median value 93%, table 2). In figure 1*b*, we further explored this scenario by adding a second contraction at time T_{pq} , corresponding to the plague and/or the earthquake, but the method could not distinguish between the three scenarios. Furthermore, the parameters estimated for the scenarios in figure 1*b* suggest that if there was a population size contraction at time T_{pq} , it was small (table 2) with median values ranging between 10% and 20%. The median values for the contraction corresponding to the abandonment of Sagalassos are very similar (90%) to the value found when there is only one major contraction (the AM scenario). Hence, the data suggest one major demographic event corresponding to the abandonment of the city but it cannot be determined whether the earthquake or the plague also contributed to changes in genetic diversity.




Figure 2. Contour maps of pairwise F_{ST} values between each of the three chronological samples of this study, Roman Sagalassos (a), Middle Byzantine Sagalassos (b), modern Ağlasun (c), and 77 modern populations from the database. Genetic distances are represented by a gradient from green (smallest F_{ST} values) to white (largest F_{ST} values). Approximate location of samples used for the analysis is indicated by a cross. Countries without any samples available are shaded. The location of Sagalassos is indicated by a star.

4. Discussion

4.1. The mtDNA pool in the Sagalassos study region across time

The maternal genetic composition of Sagalassos in Roman times shows a typical West Eurasian pool, as previously observed in the Middle Byzantine sample [14]. The multiple analyses carried out at the level of mtDNA genetic variation (F_{ST} genetic distances, PCA, shared haplotypes) reveal a clear Southwest Asian affinity, in particular with the Caucasus, Iraq, Iran, the Levant and the general Anatolian region. Genetic distances increase eastward across the Asian continent, and westward in Europe, though a genetic affinity was observed with continental Western Europe (France and Germany) and Southern Italy. Furthermore, the analyses reveal a common genetic thread along Southwest Asia, the Mediterranean basin (e.g. Southern Italian and Greek samples, including Bronze Age Crete) up to ancient central

Table 2. Demographic parameters for the scenarios described in figure 1b. The posterior weighted (ω) median, 5% and 95% percentile values are represented for the parameters described in figure 1b. The priors were taken from broad uniform distributions, ranging between a minimum and maximum value (U : min–max), for each of the parameters. The T_{pq} parameter represents the time in generations ago (assuming that one generation corresponds to 25 years and that the present is set to 2000 CE) for a possible size contraction in Sagalassos, prior to the abandonment of the city, and it has a range of 15 generations to include the plague outbreak and seventh century earthquake.

scenarios	parameters	prior	posterior		
			ω P5%	ω median	ω P95%
AM	p_a	U : 0.01–1	0.0438	0.0774	0.2211
	N_5	U : 1000–40 000	20 227	33 550	39 506
	N_{anc}	U : 1000–5000	2504	4062	4916
PQ + AM	T_{pq}	U : 50–65	50	53	63
	p_{pq}	U : 0.2–1	0.3581	0.7701	0.9777
	p_a	U : 0.01–1	0.0325	0.1086	0.5703
	N_5	U : 1000–40 000	15 413	32 032	39 750
	N_{anc}	U : 1000–5000	3219	4187	4828
PQR + AM	T_{pq}	U : 50–65	51	56	65
	p_{pq}	U : 0.2–1	0.4271	0.8939	0.9950
	p_a	U : 0.01–1	0.0371	0.0806	0.3288
	N_5	U : 1000–40 000	11 923	32 272	39 820
	N_{anc}	U : 1000–5000	2204	4109	4975

European farming communities, a pattern that may reflect the complex connections that took place in these areas across time since the Neolithic [23,24].

Roman times were certainly dynamic in the whole Mediterranean region. In that period, Sagalassos grew into the major regional centre of Pisidia. The city expanded and grounded its economic success in local agricultural and artisanal production [25]. As a dynamic centre in an empire characterized by mobility, Sagalassos may have attracted people from all over the Mediterranean and the Near East to some degree. Remarkably, the emperor Augustus established colonies of Roman veterans from Italy—with attested origins in the region of Campania—and Southern Spain in the region of Pisidia upon its incorporation into the Empire after 25 BCE [7]. Overall, such a scenario may explain the high diversity and the degree of affinity of Sagalassos with West Eurasian populations across Southwest Asia and the Mediterranean basin.

The mtDNA pool of Sagalassos and Ağlasun remained homogeneous across time, in fact no significant differences in haplogroup frequencies were observed, except for macro-haplogroup M. This is corroborated by the network analysis, which showed close genealogical relationships between most of the mtDNA lineages in the three samples.

In agreement with published data from modern Turkish populations [26], lineages of East Eurasian descent assigned to macro-haplogroup M were found in the modern sample from Ağlasun. This haplogroup is significantly more frequent in Ağlasun (15%) than in Byzantine Sagalassos, where it is absent (the non-significant value of the Roman sample might be most likely due to its low sample size), thus indicating that this East Eurasian component may have been introduced later, either recently or even as early as the Seljuk invasion of Turkey in the eleventh century CE.

Middle Byzantine Sagalassos overall showed lower genetic distances with the modern populations as opposed to the Roman sample and Ağlasun (figure 2). This could be a mere sampling effect, nevertheless the older chronological age of the Roman sample on the one hand, and isolation and drift effects in Ağlasun (as opposed to the high dynamicity of ancient Sagalassos) on the other hand, may explain a similar pattern (electronic supplementary material, text).

4.2. Demographic change

Our results support population continuity until the present in the study region of Sagalassos since Roman times, suggesting that, even though migratory events may have taken place across the centuries

(e.g. linked to the introduction of Asian lineages), no significant major changes that could not be explained by a relatively simple demographic model occurred in the mtDNA pool of the people inhabiting the area.

There is very scarce direct evidence of human presence in the immediate surroundings of Sagalassos and the Ağlasun Valley before the origins of the human communities in the later fifth century BCE. However, in all demographic scenarios tested we set the BO-phase (approx. 1000 BCE) as the initial date of human occupation of the region in light of the strong increase in cereal cultivation, arboriculture and grazing grounds for livestock, which are indicative of a significant increase of human activities in the area [27].

Our ABC analyses strongly support a scenario in which a population decline in the study region coincided with the abandonment of Sagalassos in the early thirteenth century CE, rather than with the Plague of Justinian or the earthquake in the seventh century CE (figure 1*a*).

We further explored this scenario in more depth by investigating whether the plague and/or the earthquake contributed to a reduction in population size in the region prior to the abandonment of the city. We also included a scenario of slight recovery in population size between the two contractions (figure 1*b*). Our simulations did not make it possible to discriminate with high support any of the three alternative scenarios investigated. However, cumulative posterior probabilities of the models with two contractions (71%) suggest that a decline coincident with the plague and/or the earthquake may have taken place. If it occurred, then it was moderate, with a reduction of 10–20% in effective population size in the general study region. Whether this contraction was due to an increased death rate or to migration out of the region remains to be investigated.

Most likely, either the plague and/or the earthquake in the seventh century CE determined a general loss of social complexity of Sagalassos' community [12], which would continue to wax and wane into the early thirteenth century. By then, the wider region had seen the arrival of the Seljuk Turks, who are attested to have built a caravanserai and *hamam* in Ağlasun in the 1230s CE [13]. We cannot exclude that at least part of the population of Sagalassos moved upon its abandonment, or even earlier, to Ağlasun. Nevertheless, our simulations indicate that concurrently with the abandonment of Sagalassos, a reduction of at least 90% in the effective population size of the general study region took place, suggesting that local people probably moved to more distant areas in southwest Anatolia.

As a final consideration, after accounting for the effective population size of mtDNA (one-quarter of the autosomal DNA) and the effective size as approximately one-third of the census size, the results of our simulations indicate a wide discrepancy with the census size estimated through archaeological survey—up to 50 000 including the urban and rural area (electronic supplementary material, text)—prior to any possible demographic contraction. These results should be treated with caution, owing to possible underestimations in the methodology of the archaeological survey, and to the limited power of demographic reconstructions based on a single genetic locus. In addition, the effective population sizes estimated from genetic data can be difficult to relate to actual population sizes when populations are structured. In such cases, the estimated effective size will be influenced by the area over which gene flow occurs.

The discrepancy observed might suggest that the mtDNA pool of people in the region was enriched across time by the introduction of novel lineages, thus indicating more complex scenarios and supporting the hypothesis of genetic signatures left by mobility into the area. The models tested are clearly not exhaustive of all possible demographic factors that may have taken place across time in the study region. Modelling the source and extent of gene flow in the study region of Sagalassos was beyond the scope of this study, but may become an interesting focus of future studies when additional aDNA data can be obtained across wider regions of Anatolia and the surrounding regions.

5. Conclusion

In this study, we addressed questions regarding the demographic trajectories across the last two millennia of a human population at the very fine scale of one city—Sagalassos, in southwest Turkey—and its surrounding region. By comparing mtDNA variation in three period groups, Roman, Middle Byzantine and modern, and by simulating possible scenarios inferred by historical and archaeological evidence, our data suggest that concurrently with the abandonment of the city in the early thirteenth century CE, the population of the region may have been drastically reduced by almost 90%, most likely owing to migrations towards farther regions in southwest Anatolia. Furthermore, an earlier but milder contraction in population size may have taken place in the sixth–seventh centuries CE, either owing to the Plague of Justinian and/or an earthquake in the region. Whether this hypothetical contraction was

due to increased death rate or migration out of the region remains unclear, and—together with the overall demographic scenario described here—needs to be better addressed by incorporating nuclear markers on a large genomic scale.

Data accessibility. The DNA sequences generated in this study are available in GenBank (accession number KT892706–KT892729).

Authors' contributions. C.O., R.D. and J.P. conceived the study. C.O., R.R., L.C., J.P. and R.W. designed the demographic scenarios and R.R. performed the simulations analyses. C.O. carried out the molecular laboratory work and the statistical analyses. K.V.V. collected the archaeological samples. J.C., P.T. and J.P. conducted archaeological research at Sagalassos. C.O. and R.R. drafted the manuscript. All authors revised the manuscript and gave final approval for publication.

Competing interests. We have no competing interests.

Funding. This research was supported by the Belgian Programme on Interuniversity Poles of Attraction (IAP 07/09; <http://iap-cores.be/>), the Research Fund of the University of Leuven (GOA 13/04), the KU Leuven BOF Centre of Excellence Financing on 'Centre for Archaeological Sciences 2—New methods for research in demography and interregional exchange', and Projects G.0562.11 and G. 0637.15 of the Research Foundation Flanders (FWO) (<http://sagalassos.be/en/projects>). R.R. is supported by the Leverhulme Trust (UK) under Programme grant F/00 212/AM. R.W. is supported by the ERC Advanced grant 'Empire of 2000 Cities' allotted to the Institute of History of Leiden University.

Acknowledgements. We thank Prof. Marc Waelkens, the founding director of the Sagalassos Project, for having initiated this type of interdisciplinary research and continuing to support the project, the Ministry of Culture and Tourism of the Republic of Turkey, its Kültür Varlıkları ve Müzeler Genel Müdürlüğü, its annual representatives, and the Burdur Museum for the excavation and sampling permissions, support and most appreciated aid during the fieldwork campaigns. We also thank Reyhan Yaka, METU University of Ankara, for useful assistance in the laboratory work. This research used the ALICE High Performance Computing Facility at the University of Leicester.

References

1. Brandt G *et al.* 2013 Ancient DNA reveals key stages in the formation of Central European mitochondrial genetic diversity. *Science* **342**, 257–261. (doi:10.1126/science.1241844)
2. Rasteiro R, Chikhi L. 2013 Female and male perspectives on the Neolithic transition in Europe: clues from ancient and modern genetic data. *PLoS ONE* **8**, e60944. (doi:10.1371/journal.pone.0060944)
3. Ghirotto S *et al.* 2013 Origins and evolution of the Etruscans' mtDNA. *PLoS ONE* **8**, e55519. (doi:10.1371/journal.pone.0055519)
4. Vai S *et al.* 2015 Genealogical relationships between early medieval and modern inhabitants of Piedmont. *PLoS ONE* **10**, e0116801. (doi:10.1371/journal.pone.0116801)
5. Poblome J. 2014 *Exempli gratia: Sagalassos, Marc Waelkens and interdisciplinary archaeology*. Leuven, Belgium: Leuven University Press.
6. Duru R. 2008 *From 8000 BC to 2000 BC: six thousand years of the Burdur—Antalya region*. Antalya, Turkey: Ege Yayınları.
7. Waelkens M. 2002 Romanization in the East: a case study. Sagalassos and Pisidia (SW Turkey). *IstMitt* **52**, 311–368.
8. Arndt A, Van Neer W, Hellemans B, Robben J, Volckaert F, Waelkens M. 2003 Roman trade relationships at Sagalassos (Turkey) elucidated by ancient DNA of fish remains. *J. Archaeol. Sci.* **30**, 1095–1105. (doi:10.1016/S0305-4403(02)00204-2)
9. Wagner DM *et al.* 2014 *Yersinia pestis* and the Plague of Justinian 541–543 AD: a genomic analysis. *Lancet Infect. Dis.* **14**, 319–326. (doi:10.1016/S1473-3099(13)70323-2)
10. Allen P. 1979 The Justinian Plague. *Byzantion* **49**, 5–20.
11. Waelkens M, Sintubin M, Muchez P, Paulissen E. 2000 Archaeological, geomorphological and geological evidence for a major earthquake at Sagalassos (SW Turkey) around the middle of the seventh century AD. *Geol. Soc. Lond. Spec. Publ.* **171**, 373–383. (doi:10.1144/gsl.sp.2000.171.01.27)
12. Poblome J. 2014 Shifting societal complexity in Byzantine Asia Minor and Dark Age pottery. In *LRCW 4 - archaeology and archaeometry. The Mediterranean: a market without frontiers* (eds N Poulou-Papadimitriou, E Nodarou, V Kilikoglou), pp. 623–642. Oxford, UK: Archaeopress.
13. Vanhaverbeke H, Başağaç Ö, Paul K. 2005 A Selçuk Hamam at Ağlasun (Burdur province). *Turcica* **37**, 309–336. (doi:10.2143/TURC.37.0.201709)
14. Ottoni C, Ricaut FX, Vanderheyden N, Brucato N, Waelkens M, Decorte R. 2011 Mitochondrial analysis of a Byzantine population reveals the differential impact of multiple historical events in South Anatolia. *Eur. J. Hum. Genet.* **19**, 571–576. (doi:10.1038/ejhg.2010.230)
15. Jehaes E. 1998 *Optimization of methods and procedures for the analysis of mtDNA sequences and their applications in molecular archaeological and historical finds*. Leuven, Belgium: Leuven University Press.
16. Excoffier L, Lischer HEL. 2010 Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* **10**, 564–567. (doi:10.1111/j.1755-0998.2010.02847.x)
17. Bandelt HJ, Forster P, Rohl A. 1999 Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* **16**, 37–48. (doi:10.1093/oxfordjournals.molbev.a026036)
18. Beaumont M. 2008 Joint determination of topology, divergence time and immigration in population trees. In *Simulation, genetics and human prehistory* (eds S Matsumura, P Forster, C Renfrew), pp. 135–154. Cambridge, UK: McDonald Institute for Archaeological Research.
19. Beaumont MA, Zhang W, Balding DJ. 2002 Approximate Bayesian computation in population genetics. *Genetics* **162**, 2025–2035.
20. Anderson CNK, Ramakrishnan U, Chan YL, Hadly EA. 2005 Serial SimCoal: a population genetics model for data from multiple populations and points in time. *Bioinformatics* **21**, 1733–1734. (doi:10.1093/bioinformatics/bti154)
21. Excoffier L, Novembre J, Schneider S. 2000 SIMCOAL: a general coalescent program for the simulation of molecular data in interconnected populations with arbitrary demography. *J. Hered.* **91**, 506–509. (doi:10.1093/jhered/91.6.506)
22. Csilléry K, François O, Blum MGB. 2012 abc: an R package for approximate Bayesian computation (ABC). *Methods Ecol. Evol.* **3**, 475–479. (doi:10.1111/j.2041-210X.2011.00179.x)
23. Paschou P *et al.* 2014 Maritime route of colonization of Europe. *Proc. Natl Acad. Sci. USA* **111**, 9211–9216. (doi:10.1073/pnas.1320811111)
24. Hughey JR, Paschou P, Drineas P, Mastropalo D, Lotakis DM, Navas PA, Michalodimitrakis M, Stamatoyannopoulos JA, Stamatoyannopoulos G. 2013 A European population in Minoan Bronze Age Crete. *Nat Commun.* **4**, 1861. (doi:10.1038/ncomms2871)
25. Poblome J, Vienne W, Kucha H, Waelkens M, Laduron D, Depuydt F. 1997 The clay raw materials of Sagalassos Red Slip ware - a chronological evaluation. In *Sagalassos IV* (eds M Waelkens, J Poblome), pp. 507–518. Leuven, Belgium: Leuven University Press.
26. Quintana-Murci L *et al.* 2004 Where west meets east: the complex mtDNA landscape of the southwest and Central Asian corridor. *Am. J. Hum. Genet.* **74**, 827–845. (doi:10.1086/383236)
27. Vermeore M. 2004 *Holocene vegetation history in the territory of Sagalassos, southwest Turkey a palynological approach*. Turnhout, Belgium: Brepols.