

Supplementary Information

Quantitative proteomics screen identifies a substrate repertoire of rhomboid protease RHBDL2 in human cells and implicates it in epithelial homeostasis.

¹Nicholas Johnson, ^{1,2}Jana Březinová, ³Elaine Stephens, ⁵Emma Burbridge, ^{3,4}Matthew Freeman, ⁵Colin Adrain* and ¹Kvido Strisovsky*

¹Institute of Organic Chemistry and Biochemistry, Czech Academy of Science, Flemingovo n. 2, Prague, 166 10, Czech Republic

²Department of Biochemistry, Faculty of Science, Charles University, Prague, Czech Republic

³MRC Laboratory of Molecular Biology, Cambridge, CB2 2QH, United Kingdom

⁴Sir William Dunn School of Pathology, Oxford, OX1 3RE, United Kingdom

⁵Instituto Gulbenkian de Ciência, Lisbon, Portugal

*Authors for correspondence (email: kvido.strisovsky@uochb.cas.cz, cadrain@igc.gulbenkian.pt)

Table S1: The full dataset of proteins identified in the SILAC experiment.

The table summarizes all proteins with at least one quantifiable SILAC pair identified by MaxQuant¹⁵ with corresponding topology predictions by Phobius¹⁷.



Table S1 rev1.xlsx

Please open attachment from the sidebar.

Dataset S1: Results of proteomics experiment to identify RHBDL2 substrates (Proteator format).



Proteator results
Johnson et al.html

Please open attachment from the sidebar.